



Hyperbolization of cusps with convex boundary

François Fillastre, Ivan Izmistiev, Giona Veronelli

► To cite this version:

François Fillastre, Ivan Izmistiev, Giona Veronelli. Hyperbolization of cusps with convex boundary. 2015. hal-01152466

HAL Id: hal-01152466

<https://hal.science/hal-01152466>

Preprint submitted on 17 May 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HYPERBOLIZATION OF CUSPS WITH CONVEX BOUNDARY

FRANÇOIS FILLASTRE, IVAN IZMESTIEV, AND GIONA VERONELLI

ABSTRACT. We prove that for every metric on the torus with curvature bounded from below by -1 in the sense of Alexandrov there exists a hyperbolic cusp with convex boundary such that the induced metric on the boundary is the given metric. The proof is by polyhedral approximation.

This was the last open case of a general theorem: every metric with curvature bounded from below on a compact surface is isometric to a convex surface in a 3-dimensional space form.

CONTENTS

1. Introduction	1
1.1. Statement of the results	1
1.2. Hyperbolization of products manifolds	3
1.3. Smooth variational approach?	5
1.4. Acknowledgement	6
2. Background	6
2.1. CBB metrics on compact surfaces	6
2.2. Horoconvex functions	7
2.3. Induced metric	10
3. Proof of Theorem 1.1	11
3.1. A uniform bound on horographs	12
3.2. Convergence of groups	12
3.3. Construction of the solution	13
3.4. Convergence of metrics	13
References	16

1. INTRODUCTION

1.1. Statement of the results. Let T denote the 2-dimensional torus. A hyperbolic cusp C with convex boundary is a complete hyperbolic manifold of finite volume, homeomorphic to $T \times [0, +\infty[$, and such that the boundary $\partial C = T \times \{0\}$ is geodesically convex. By the Buyalo convex hypersurface theorem [AKP08], the induced inner metric on ∂C has curvature bounded from below by -1 in the sense of Alexandrov — in short, the metric is CBB(-1), see Section 2.1. In the present paper, we prove that all the CBB(-1) metrics on the torus are obtained in this way.

Date: May 15, 2015

v1.

Theorem 1.1. *Let m be a $\text{CBB}(-1)$ metric on the torus. Then there exists a hyperbolic cusp C with convex boundary such that the induced metric on ∂C is isometric to m .*

Throughout the paper, by induced metric we mean an intrinsic metric. Examples of $\text{CBB}(-1)$ metrics on the torus are distances defined by Riemannian metrics of curvature ≥ -1 . Using classical regularity results of Pogorelov (see below), Theorem 1.1 implies the following.

Theorem 1.2. *Let g be a smooth Riemannian metric of curvature > -1 on the torus. Then there exists a hyperbolic cusp C with smooth strictly convex boundary such that the induced Riemannian metric on ∂C is isometric to g .*

Another examples of $\text{CBB}(-1)$ metrics on the torus are hyperbolic metrics with conical singularities of positive curvatures. Recall that the (singular) curvature at a cone singularity is 2π minus the total angle around the singularity. The proof of Theorem 1.1 will be done by polyhedral approximation, using the following result.

Theorem 1.3 ([FI09]). *Let m be a hyperbolic metric with conical singularities of positive curvature on the torus. Then there exists a hyperbolic cusp C with convex polyhedral boundary such that the induced inner metric on ∂C is isometric to m .*

Actually it is proved in [FI09] that the cusp C in the statement of Theorem 1.3 is unique. One can hope a uniqueness result in Theorem 1.1 (that would also imply uniqueness in Theorem 1.2). This should be the subject of a forthcoming paper.

The above statements were the last steps in order to get the following general statement.

Theorem 1.4. *Let m be a $\text{CBB}(k)$ metric on a compact surface. Then m is isometric to a convex surface S in a Riemannian space of constant curvature k . Moreover*

- *if m is a metric of curvature k with conical singularities of positive curvature, then S is polyhedral,*
- *if m comes from a smooth Riemannian metric with curvature $> k$, then S is smooth and strictly convex.*

In the rest of the introduction we will recall all the results leading to Theorem 1.4. The proof of Theorem 1.1 is based on a general result of polyhedral approximation that is recalled in Section 2.1 (Theorem 2.2). Going to the universal cover, boundaries of convex hyperbolic cusps are seen as convex surfaces of the hyperbolic space invariant under the action of a group of isometries acting cocompactly on a horosphere. Such surfaces are graphs of *horconvex functions* defined on the horosphere. They are introduced in Section 2.2. They can be written in terms of convex functions on the plane, hence they will inherit strong properties from convex functions.

Then we prove Theorem 1.1 in Section 3. Theorem 1.3 and Theorem 2.2 give a sequence of polyhedral surfaces in \mathbb{H}^3 . One shows that this sequence converges to a convex surface, invariant under the action of a group Γ . The main point is to check that the induced metric on the quotient of the surface by Γ is isometric to the metric m (Section 3.4).

1.2. Hyperbolization of products manifolds. The notion of metric space with non-negative curvature was introduced by A.D. Alexandrov in order to describe the induced metric on the boundary of convex bodies in \mathbb{R}^3 . He proved that this property characterizes the convex bodies in the following sense.

Theorem 1.5 ([Ale06]). *Any CBB(0) metric on the sphere is isometric to the boundary of a convex body in \mathbb{R}^3 .*

The hyperbolic version is as follows.

Theorem 1.6 ([Ale06]). *Any CBB(-1) metric on the sphere is isometric to the boundary of a compact convex set in \mathbb{H}^3 .*

Let us mention the following general local result, obtained from Theorem 1.6 and a gluing theorem.

Theorem 1.7 ([Ale06]). *Each point in a CBB(-1) surface has a neighbourhood isometric to a convex surface in \mathbb{H}^3 .*

Forgetting the part of the hyperbolic space outside of the convex body, one derives from Theorem 1.6 the following hyperbolization theorem for the ball.

Theorem 1.8. *Let m be a CBB(-1) metric on the sphere. Then there exists a hyperbolic ball M with convex boundary such that the induced metric on ∂M is isometric to m .*

Actually Theorem 1.8 also implies Theorem 1.6, because in this case the developing map is an isometric embedding [CEG06, Proposition I.1.4.2.]. Theorem 1.6 and Theorem 1.8 are proved by polyhedral approximation. For example Theorem 1.6 is proved from the following particular case.

Theorem 1.9. *Let m be a hyperbolic metric with conical singularities of positive curvature on the sphere. Then there exists a hyperbolic ball M with convex polyhedral boundary such that the induced metric on ∂M is isometric to m .*

The following regularity result roughly says that if the metric on a convex surface in \mathbb{H}^3 is smooth, then the surface is smooth.

Theorem 1.10 ([Pog73, Theorem 1, chap. V 8]). *Let S be a surface with a C^k , $k \geq 5$, Riemannian metric of curvature > -1 . If S admits a convex isometric embedding into \mathbb{H}^3 , then its image is a C^{k-1} surface.*

See [CX15] for more precise results if S is homeomorphic to the sphere, in particular if the curvature is ≥ -1 . Theorem 1.10 and Theorem 1.6 immediately give the following.

Theorem 1.11. *Let g be a smooth Riemannian metric with curvature > -1 on a the sphere. Then there exists a hyperbolic ball M with smooth convex boundary such that the induced metric on ∂M is isometric to g .*

For metrics on a compact (connected) surface S of genus > 1 , the following result was recently proved.

Theorem 1.12 ([Slu, Slu14]). *Let M be a compact connected 3-manifold with boundary of the type $S \times [1, 1]$. Let m be a $CBB(-1)$ metric on ∂M . Then there exists a hyperbolic metric in M with a convex boundary such that the induced metric on ∂M is isometric to m .*

The proof of Theorem 1.12 goes by smooth approximation. The smooth version of Theorem 1.12 is included in the following more general result.

Theorem 1.13 ([Lab92]). *Let M be a compact manifold with boundary (different from the solid torus) which admits a structure of a strictly convex hyperbolic manifold. Let g be a smooth metric on ∂M of curvature > -1 . Then there exists a convex hyperbolic metric on M which induces g on ∂M .*

See also [Sch06], which contains a uniqueness result. Of course one can take for metrics m in the statement of Theorem 1.12 hyperbolic metrics on S with conical singularities with positive curvature. But the boundary of the solution is not necessarily of polyhedral type. This is because the boundary may meet the boundary of the convex core of M . If M is *Fuchsian*, that is if its convex core is a totally geodesic surface, this cannot happen. Actually we have the following.

Theorem 1.14 ([Fil07]). *Let m be a hyperbolic metric with conical singularities with positive curvature on a compact surface S of genus > 1 . Then there exists a Fuchsian hyperbolic manifold $S \times [-1, 1]$ with polyhedral convex boundary such that the induced metric on the boundary components is isometric to m .*

The smooth analogue was known for a long:

Theorem 1.15 ([Gro86]). *Let g be a smooth Riemannian metric with curvature > -1 on a compact surface S of genus > 1 . Then there exists a Fuchsian hyperbolic manifold $S \times [-1, 1]$ with smooth convex boundary such that the induced metric on the boundary components is isometric to g .*

Using Theorem 1.15 instead of Theorem 1.13, the proof of Theorem 1.12 leads to the following.

Theorem 1.16. *Let m be a $CBB(-1)$ metric on a compact surface S of genus > 1 . Then there exists a Fuchsian hyperbolic manifold $S \times [-1, 1]$ with convex boundary such that the induced metric on the boundary components is isometric to m .*

Let us put all these statements together. Cutting in a suitable way the hyperbolic manifolds given in theorems 1.6, 1.11, 1.9, 1.1, 1.2, 1.3, 1.16, 1.15, 1.14, we obtain the following result.

Theorem 1.17. *Let m be a $CBB(-1)$ metric on a compact surface S . Then the manifold $S \times [-1, 1]$ admits a hyperbolic metric, such that $S \times \{-1\}$ is convex and isometric to m for the induced inner metric, and $S \times \{1\}$ has constant curvature.*

Moreover

- if m is a hyperbolic metric with conical singularities of positive curvature, then $S \times \{-1\}$ is polyhedral,

- if m comes from a smooth Riemannian metric with curvature > -1 , then $S \times \{-1\}$ is smooth and strictly convex.

Note that in the case of genus > 1 , we have chosen the Fuchsian solution, but the quasi-Fuchsian Theorem 1.12 gives many choices for the realization of the prescribed metric. Actually all the cases in Theorem 1.17 share the same property: the holonomy of their developing map fixes a point (the point may not be in the hyperbolic space, see for example the beginning of Section 3).

In the case of $\text{CBB}(-1)$ metrics on the torus, we could also consider a hyperbolic metric with convex boundary on a full torus. In this direction, only the smooth case is known.

Theorem 1.18 ([Sch06]). *Let g be a smooth Riemannian metric of curvature > -1 on the torus T . Then there exists a (unique) hyperbolic metric on the full torus such that the metric on the boundary is smooth, strictly convex and isometric to g .*

Another question is to realize $\text{CBB}(-1)$ metrics on compact surfaces of genus > 1 as the convex boundary of more general compact hyperbolic manifold, analogously to Theorem 1.13.

We cited Theorem 1.5 about realization of $\text{CBB}(0)$ metrics on the sphere in the Euclidean space. There is also an analogue result about realization of $\text{CBB}(1)$ metrics on the sphere in the 3 dimensional sphere [Ale06], as well as the polyhedral and smooth counterparts. Theorem 1.17 gives all the possibilities for a $\text{CBB}(-1)$ metric on a compact surface of genus > 1 . Moreover it is obvious that any flat metric on a torus T can be extended to a flat metric on $T \times [-1, 1]$. Lemma 2.1 says that we have exhausted all the possibilities. Theorem 1.4 follows.

A question is to know if the constant curvature metric on $S \times [-1, 1]$ is unique. Due to the work of Pogorelov, the answer is positive if S is the sphere [Pog73]. As we already mentioned, in the torus case this is work in progress. As the only unsolved case there would remain that of Fuchsian hyperbolic manifolds with convex boundary.

1.3. Smooth variational approach? As we said, Theorem 1.5 was proved by polyhedral approximation. It is based on the following seminal theorem, proved in the 1940's.

Theorem 1.19 ([Ale06]). *Let m be a flat metric with conical singularities of positive curvature on the sphere. Then there exists a convex polyhedron in Euclidean space with inner induced metric m on the boundary.*

The proof of Theorem 1.19 is done by a continuity method, based on topological arguments, in particular the Domain Invariance theorem. Some years ago, a variational proof of Theorem 1.19 was given in [BI08]. The functional is a discrete Hilbert–Einstein functional. It was then used in [Izm08], and later in [FI09] to prove Theorem 1.3. A long-standing question is to use the smooth Hilbert–Einstein functional to give a variational proof of the smooth version of Theorem 1.19 (known as Weyl problem) [BH37, Izm13]. It would be interesting to give a variational proof of Theorem 1.2 as

well. There are reasons to think that the functional will have good properties in the case of a hyperbolic cusp.

1.4. Acknowledgement. The authors thank Stephanie Alexander, Thomas Richard, Joël Rouyer, Dima Slutskiy for useful conversations.

The first author was partially supported by the ANR GR-Analysis-Geometry and by the mathematic department of the Universidade Federal do Rio de Janeiro and this research has been conducted as part of the project Labex MME-DII (ANR11-LBX-0023-01). A part of this work was done during his stay at the Universidade Federal do Rio de Janeiro. He thanks this institution for its hospitality.

The second author was supported by the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC Grant agreement no. 247029-SDModels.

The third author was partially supported by the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA).

2. BACKGROUND

2.1. CBB metrics on compact surfaces. We follow [BBI01] for basic definitions and results about metric geometry. See also [BH99] and [AKP]. Let m be a metric on a compact surface S (by this we imply that the topology given by m is the topology of S). We suppose that m is intrinsic, that is for any $x, y \in S$, $m(x, y)$ is equal to the infimum of the length of the continuous curves between x and y . By the Hopf–Rinow theorem, there always exists a shortest path between x and y .

The metric m is $\text{CBB}(k)$ if every point has a neighbourhood U such that any triangle contained in U is thicker than the comparison triangle in the model space of constant curvature k (see the references above for precise and equivalent definitions). By the Toponogov globalization theorem, this property is actually true for any triangle in (S, m) .

A shortest path between two points in a $\text{CBB}(k)$ space may not be unique, as show the example of a disc of curvature k with a sector removed and the two resulting sides identified. But shortest paths in $\text{CBB}(k)$ do not branch.

Let (S, m) be a polyhedral $\text{CBB}(k)$ metric, that is a metric of constant curvature k with singular curvatures k_i (the k_i have to be positive [BBI01, 10.9.5]). It has to satisfy the Gauss–Bonnet formula

$$2\pi\chi(S) = k \text{ area}(S) + \sum k_i$$

i.e.

$$2\pi\chi(S) \geq k \text{ area}(S)$$

with equality if and only if m is a smooth constant curvature metric (no conical singularities).

Now let (S, m) be any $\text{CBB}(k)$ metric. By a theorem of Alexandrov and Zalgaller, (S, m) can be decomposed into non-overlapping geodesic triangles. Replacing each triangle by a comparison triangle in the space of constant curvature k , we obtain a polyhedral $\text{CBB}(k)$ metric on S see [Ric12, IRV15] for details. In particular, we obtain the following.

Lemma 2.1. *A compact surface S can be endowed with a $CBB(k)$ metric if and only if*

- *if S is a sphere, $k \in \mathbb{R}$,*
- *if S is a torus, $k = 0$ and the metric is a flat Riemannian metric or $k < 0$,*
- *otherwise, $k < 0$.*

Actually, Alexandrov and Zalgaller proved much more, but in a different context. Roughly speaking, the triangulation of the $CBB(-1)$ can be chosen as fine as wanted. If the perimeter of the triangles goes to 0, then the sequence of polyhedral metrics obtained by replacing the triangles by comparison triangles converge to (S, m) in the Gromov–Hausdorff sense [Ric12, IRV15].

Theorem 2.2. *Let d be a $CBB(-1)$ metric on a compact surface. Then there exists polyhedral $CBB(-1)$ sequence of metric m_n on the torus Gromov Hausdorff converging to m .*

At the end of the day, in the case of the torus, it is not hard to conclude from Theorem 1.1 and Proposition 3.5 that the convergence can be taken uniform in Theorem 2.2.

Let us mention the following results about Gromov–Hausdorff convergence that we will use in the sequel.

Lemma 2.3 ([BH99, I.5.40], [BBI01, 7.3.14]). *Gromov–Hausdorff convergence implies convergence of the diameter*

Theorem 2.4 ([BGP92], [BBI01, 10.10.11]). *If a sequence (S, m_n) of $CBB(k)$ metrics on a compact surface S converges in the Gromov–Hausdorff sense to a $CBB(k)$ metric on S , then the sequence of the areas of (S, m_n) (the total two dimensional Hausdorff measure) is bounded from below by a positive constant.*

2.2. Horoconvex functions. We identify \mathbb{R}^2 with a given horosphere $H \subset \mathbb{H}^3$, with center ∞ . We get coordinates (x, t) on $\mathbb{H}^3 = H \times \mathbb{R}$, where t is the signed distance from a point to H : it is positive if and only if the point is in the exterior of the horoball bounded by H . Note that it is the length of the segment between x and its orthogonal projection onto H , and that the line from x to ∞ is orthogonal to H .

Let $u : \mathbb{R}^2 \rightarrow \mathbb{R}$. The *horograph* of u is the subset $(x, u(x)) \in \mathbb{H}^3$ for those coordinates. The horograph is said to be convex if the surface is convex in \mathbb{H}^3 in the sense that it bounds a geodesically convex set. In the Klein projective model, this corresponds to the affine notion of convexity.

Remark 2.5. In the upper half plane model, if the horosphere H is the horizontal plane at height 1, then the horograph of u is the graph of e^{-u} .

We have the following characterization of horographs. It was already known in the smooth case [GSS09]. Let us also mention that the Darboux equation related to Theorem 1.2 is studied in [RS94].

Proposition 2.6. *The horograph of $u : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a convex surface if and only if the function*

$$x \mapsto e^{-2u(x)} + \|x\|^2$$

is convex, with $\|\cdot\|$ the Euclidean norm.

In particular, e^{-2u} is semi-convex, or lower- C^∞ , compare with 10.33 and 13.27 in [RW98]. We will call *horoconvex* a function satisfying the hypothesis of the proposition.

Proof. As above, consider coordinates $(x, t)_H$ on $\mathbb{H}^3 = H \times \mathbb{R}$. The horograph of u is convex in \mathbb{H}^3 if and only if at each point $(x_0, u(x_0))_H$ there exists a totally geodesic surface Σ containing the point $(x_0, u(x_0))_H$ and such that $\Sigma \subset \{(x, y)_H : y \geq u(x)\}$. Now, consider on the Poincaré halfspace model the standard Euclidean coordinates $(y, s)_E \in \mathbb{R}^2 \times (0, \infty)$. Without loss of generality we can assume that the horosphere H is the plane at height 1 in this model. Then we have $(x, t)_H = (x, e^{-t})_E$. In this system, Σ has to be a half-sphere with center $(c, 0)_E$ on the plane at infinity $\mathbb{R}^2 \times \{0\}$. In particular every such a half-sphere containing the point $(x_0, u(x_0))_H$ is given by

$$\{(x, s)_E : e^{-2u(x_0)} + \|c - x_0\|^2 = \|x - c\|^2 + s^2\}.$$

Coming back to $(\cdot, \cdot)_H$ coordinates, we have obtained that the horograph of u is convex if and only if for all $x_0 \in \mathbb{R}^2$ there exists a point $c \in \mathbb{R}^2$ such that

$$u(x) \leq -\frac{1}{2} \ln \left(e^{-2u(x_0)} + \|c - x_0\|^2 - \|c - x\|^2 \right),$$

that is,

$$(1) \quad e^{-2u(x)} - e^{-2u(x_0)} \geq \|c - x_0\|^2 - \|c - x\|^2 = -\|x - x_0\|^2 + 2 \langle G, x - x_0 \rangle,$$

where $G := c - x_0 \in \mathbb{R}^2$. Since $x \in \mathbb{R}^2$ is arbitrary, (1) is equivalent to

$$(2) \quad e^{-2u(x_0+v)} + \|x_0 + v\|^2 - e^{-2u(x_0)} - \|x_0\|^2 \geq \langle 2(x_0 + G), v \rangle, \quad \forall v \in \mathbb{R}^2.$$

In turn, (2) means that if $G \in \mathbb{R}^2$ is such that (1) is satisfied, then at each point $x_0 \in \mathbb{R}^2$ the graph of the function $e^{-2u(x)} + \|x\|^2$ has the planar graph of $v \mapsto \langle 2(x_0 + G), v \rangle + e^{-2u(x_0)} + \|x_0\|^2$ as a support plane. Hence $e^{-2u(x)} + \|x\|^2$ is a convex function on \mathbb{R}^2 if and only if the horograph of u is convex in \mathbb{H}^3 . \square

Remark 2.7. Suppose that $x = x_0 + th$ for some unitary vector $h \in \mathbb{R}^2$ and $t > 0$. Then (1) reads

$$(3) \quad \frac{1}{t} \left(e^{-2u(x_0+th)} - e^{-2u(x_0)} \right) \geq -t|h|^2 + 2 \langle G, h \rangle.$$

If $u \in C^1(\mathbb{R}^2)$, taking the limit as $t \rightarrow 0$ we get that

$$\langle \text{grad}_{x_0} e^{-2u}, h \rangle \geq \langle 2G, h \rangle,$$

and since the latter inequality holds for both h and $-h$ we have necessarily that G is unique and

$$G = \frac{1}{2} \text{grad}_{x_0} e^{-2u} = -e^{-2u(x_0)} \text{grad}_{x_0} u.$$

Corollary 2.8. *Let u be horoconvex and $\epsilon > 0$. Then $u + \epsilon$ is horoconvex.*

Proof. By Proposition 2.6, one has to see that $e^{-2\epsilon} e^{-2u(x)} + \|x\|^2$ is convex, that is equivalent to the convexity of $e^{-2u(x)} + e^{2\epsilon} \|x\|^2$. Let $f + g$ be convex, g convex and $\lambda > 1$. Then $f + \lambda g = (f + g) + (\lambda - 1)g$ is convex as a sum of two convex functions. \square

Example 2.9. In one dimension, the function $t \mapsto \cos(t)/20$ is horoconvex, see Figure 1. More generally, any function C^2 close to a constant function is horoconvex. But $t \mapsto \cos(t)$ is not horoconvex. This example shows that u horoconvex does not imply λu horoconvex.




FIGURE 1. Graphs of $t \mapsto e^{-\cos(t)/10}$ and $t \mapsto e^{-2\cos(t)}$.

Horoconvex functions inherit strong properties from convex functions.

Corollary 2.10. *Any sequence of uniformly bounded horoconvex functions is equi-Lipschitz on any compact set of \mathbb{R}^2 .*

Moreover, up to extracting a subsequence, the sequence converges uniformly on any compact set to a horoconvex function.

Proof. Let $C \subset \mathbb{R}^2$ be a compact set and (u_n) a sequence of uniformly bounded horoconvex functions. Let $F_n(x) = e^{-2u_n(x)} + \|x\|^2$, which is convex by Proposition 2.6. By [Roc97, 10.4], there exists an $\epsilon > 0$ such that for any $x, y \in C$

$$|F_n(x) - F_n(y)| \leq \frac{\max_C F_n - \min_C F_n}{\epsilon} \|x - y\|.$$

As the u_n are uniformly bounded, F_n are uniformly bounded on C , hence there exists a number A satisfying

$$|F_n(x) - F_n(y)| \leq A \|x - y\|.$$

Using again that the u_n are uniformly bounded, and that

$$|\|x\|^2 - \|y\|^2| \leq 2 \left(\sup_C \|x\| \right) \|x - y\|,$$

we thus obtain that

$$|u_n(x) - u_n(y)| \leq A' \|x - y\|$$

for some $A' = A'(C) > 0$ independent of n . So the sequence is equi-Lipschitz on C .

Up to extract a subsequence, the sequence of convex functions (F_n) converge (uniformly on compact sets) to a convex function F [Roc97, Theorem 10.9]). As the u_n are uniformly bounded, there exists a positive constant c such that $F_n \geq c + \|\cdot\|^2$, so $F > \|\cdot\|^2$, hence the function

$$u = -\frac{1}{2} \ln(F - \|\cdot\|^2)$$

is well defined and horoconvex by definition. As (F_n) converges to F uniformly on compact sets, it follows easily that u_n converges to u uniformly on compact sets. \square

In particular, a horoconvex function is Lipschitz on any compact set. By Rademacher theorem, it is differentiable almost everywhere.

2.3. Induced metric. The length of a curve c in \mathbb{H}^3 is the supremum of the length of all the broken lines of \mathbb{H}^3 with vertices on c . Equivalently [BBI01, Theorem 2.7.6], if c is Lipschitz (in \mathbb{H}^3), then

$$L(c) = \int \|c'\|_{\mathbb{H}^3} .$$

Let S be a convex surface in \mathbb{H}^3 . The (intrinsic) *induced metric* on S between two points a, b of S is the infimum of all the Lipschitz curves of finite length between a and b .

Let u be a horoconvex function. Let \tilde{d}_u be the intrinsic metric induced on the horograph of u . For simplicity, we will look at metrics onto \mathbb{R}^2 rather than on the surfaces: for $(x, y) \in \mathbb{R}^2$,

$$d_u(x, y) := \tilde{d}_u \left(\left(x, e^{-u(x)} \right), \left(y, e^{-u(y)} \right) \right) .$$

Note that, in the upper half space model, notions of locally Lipschitz are equivalent for the metric of \mathbb{H}^3 and \mathbb{R}^3 . As the projection onto the horizontal plane is contracting, a locally Lipschitz curve of \mathbb{H}^3 is projected onto a locally Lipschitz curve of \mathbb{R}^2 .

Let $c : [a, b] \rightarrow \mathbb{R}^2$ be a Lipschitz curve. Let c_u be the corresponding curve in the horograph of u i.e. on the graph of e^{-u} :

$$c_u = \begin{pmatrix} c \\ e^{-u \circ c} \end{pmatrix} .$$

As u is Lipschitz, c_u is a Lipschitz curve of \mathbb{H}^3 .

Let us denote by $L_u(c)$ the length of c_u for the metric \tilde{d}_u on S_u . Using the half-space model metric

$$L_u(c) = \int_a^b \frac{\|c'_u\|_{\mathbb{R}^3}}{(c_u)_3} = \int_a^b e^{u \circ c} \left(\|c'\|^2 + Du(c')^2 e^{-2(u \circ c)} \right)^{1/2}$$

i.e.

$$(4) \quad L_u(c) = \int_a^b \left(e^{2u \circ c} \|c'\|^2 + Du(c')^2 \right)^{1/2} .$$

Lemma 2.11. *Let (u_n) be a uniformly bounded sequence of horoconvex functions. Then on any compact set K , d_{u_n} are uniformly Lipschitz equivalent to the Euclidean metric: $\exists \lambda_1, \lambda_2 > 0$ such that*

$$\lambda_1 d_{\mathbb{R}^2} \leq d_{u_n} \leq \lambda_2 d_{\mathbb{R}^2} .$$

Moreover, for any Lipschitz curve c contained in K ,

$$\lambda_1 L_{\mathbb{R}^2}(c) \leq L_{u_n}(c) \leq \lambda_2 L_{\mathbb{R}^2}(c) .$$

Proof. By construction, for any $x, y \in K$, $d_{u_n}(x, y)$ is not less than the distance in \mathbb{H}^3 between the corresponding points on the horograph of u . As K is compact and the u_n uniformly bounded, the corresponding horographs above K in \mathbb{R}^3 are contained in a hyperbolic ball. There exists a constant λ_1 such that on this ball, $d_{\mathbb{H}^3} \geq \lambda_1 d_{\mathbb{R}^3}$, and clearly $d_{\mathbb{R}^3} \geq d_{\mathbb{R}^2}$. As the length of a Lipschitz curve for d_{u_n} is the supremum of the length of broken shortest paths [BBI01, 2.3.4, 2.4.3] it follows that for any curve c in K , $\lambda_1 L_{\mathbb{R}^2}(c) \leq L_{u_n}(c)$.

On the other hand, for any curve c in K , it follows from (4)

$$L_{u_n}(c) \leq \int_a^b e^{-u_n} \|c'\| + |Du_n(c')|$$

and as the u_n are uniformly bounded on K by assumption, as well as $\|\text{grad } u_n\|$ when it exists by Corollary 2.10, then there exists λ_2 such that

$$L_{u_n}(c) \leq \lambda_2 L_{\mathbb{R}^2}(c) .$$

□

The proof of the following lemma mimics the one for convex bodies in Euclidean space [BBI01, p. 358].

Lemma 2.12. *Let u, v be horoconvex such that $u \leq v$ and*

$$\delta = \sup_{\mathbb{R}^2}(v - u)$$

is finite. Then

$$d_u \leq d_v + 2\delta .$$

Proof. Recall that the Hyperbolic Busemann–Feller Lemma, [BH99, II.2.4] says that the orthogonal projection onto a convex set in \mathbb{H}^3 is contracting. $u \leq v$ implies that the horograph of v is in the exterior of the convex set bounded by the horograph of u , so the orthogonal projection is well defined from the horograph of v onto the horograph of u . For $a \in \mathbb{R}^2$, let $p_\perp(a)$ be the vertical projection onto \mathbb{R}^2 of the orthogonal projection of $(a, v(a))$ onto the horograph of u .

On one hand, Busemann–Feller Lemma implies that

$$(5) \quad d_u(p_\perp(a), p_\perp(b)) \leq d_v(a, b) .$$

On the other hand, Busemann–Feller Lemma implies that $d_u(p_\perp(a), a)$ is less than the distance in \mathbb{H}^3 between $(a, v(a))$ and $(a, u(a))$, and this last quantity is less than δ by assumption, so

$$(6) \quad d_u(p_\perp(a), a) \leq \delta .$$

The result follows from (5), (6) and the triangle inequality. □

3. PROOF OF THEOREM 1.1

Now let (S, m) be a CBB(−1) metric on the torus. According to Theorem 2.2 there exists a sequence of polyhedral CBB(−1) metric m_n on the torus Gromov–Hausdorff converging to m . By Theorem 1.3, for any n there exists a hyperbolic cusp C_n with convex boundary and induced metric on ∂C_n isometric to m_n .

As mentioned in the introduction, the universal cover \widetilde{C}_n can be isometrically embedded as a convex subset of \mathbb{H}^3 via the developing map D . The action of the fundamental group $\pi_1(C_n) \cong \pi_1(T)$ on \widetilde{C}_n by deck transformations yields a representation $\rho : \pi_1(T) \rightarrow \text{Iso}^+(\mathbb{H}^3)$. The cusp C_n contains a totally umbilic torus M with Euclidean metric. It follows that the developing map maps the universal cover of M to a horosphere. The group $\rho(\pi_1(C_n)) = \Gamma_n$ acts on $D(\widetilde{M})$ freely with a compact orbit space. The group Γ_n is a group of parabolic isometries. The surface $S_n = \partial D(\widetilde{C}_n)$ is convex and globally invariant under the action of Γ_n . It is easy to see that S_n is homeomorphic to H via the central projection from the center of $D(\widetilde{M})$. Up to rotations of the hyperbolic space, we normalize the surfaces S_n in such a way that the point fixed by Γ_n is ∞ in the half space model. Moreover, choosing a point x_0 in the universal cover of the torus, up to compose by parabolic and hyperbolic isometries, we consider that the developing maps send x_0 onto $(0, 0, 1)_E$.

So the surfaces S_n are described by horoconvex functions u_n . We identify Γ_n with the corresponding lattice in $H = \mathbb{R}^2$. In particular, $u_n(\gamma \cdot x) = u_n(x)$ for any $\gamma \in \Gamma_n$ and $x \in \mathbb{R}^2$, and the normalization above says that $u_n(0) = 0$ for any n . The quotient of d_{u_n} by Γ_n is isometric to m_n .

We will prove that (u_n, Γ_n) converge to some (u, Γ) and that the quotient of d_u by Γ is isometric to m . This will prove Theorem 1.1: the wanted cusp is the quotient by Γ of the convex side of the horograph of u .

3.1. A uniform bound on horographs. From Lemma 2.3, there exists a uniform upper bound diam of all the diameters of the metrics m_n .

Lemma 3.1. (1) *The sequence $(u_n)_n$ is uniformly bounded.*

(2) *There is a compact set $D \subset \mathbb{R}^2$ such that for any $y \in \mathbb{R}^2$ and any n , there exists $\gamma \in \Gamma_n$ with $\gamma \cdot y \in D$.*

Proof. Recall that all the horographs S_n of u_n pass through $x_0 = (0, 0, 1)_E$. Let b_n be the set of points on S_n at distance $\leq \text{diam}$ from $(0, 0, 1)_E$. Then as the distance on S_n is greater than the extrinsic distance of \mathbb{H}^3 , all the b_n are contained in a same hyperbolic ball B .

In the half space model, let D be the projection of the ball B onto the horizontal plane passing through the origin. Observe that $D = \overline{B_\delta^{\mathbb{R}^2}(0)}$ is a Euclidean closed ball centred at the origin of \mathbb{R}^2 . As B is contained between two horosphere centred at ∞ (i.e. two horizontal plane) the u_n are uniformly bounded on D , say $c_1 < u_n < c_2$. Now by construction, for any $y \in \mathbb{R}^2$, there exists $\gamma \in \Gamma_n$ such that $\gamma \cdot y \in D$. Hence $c_1 < u_n(\gamma \cdot y) = u_n(y) < c_2$. \square

By Lemma 3.1 (1) and Corollary 2.10, we get

Corollary 3.2. *The sequence $(u_n)_n$ is equi-Lipschitz on every compact set of \mathbb{R}^2 .*

3.2. Convergence of groups.

Lemma 3.3. *There exists a sequence $(a_n, b_n)_n$ of generators of Γ_n which converge in \mathbb{R}^2 (up to extract a subsequence) to two linearly independent non-zero vectors a and b .*

Proof. Let us choose generators (a_n, b_n) of Γ_n which are contained in $\overline{B_{3\delta}^{\mathbb{R}^2}(0)}$ (here we identify an element of γ of Γ_n with the vector $\gamma \cdot 0$ of \mathbb{R}^2), that is possible by Lemma 3.1. Since $\overline{B_{3\delta}^{\mathbb{R}^2}(0)}$ is compact, up to take a subsequence we get the existence of two vectors $a, b \in \mathbb{R}^2$ such that $a_n \rightarrow a$ and $b_n \rightarrow b$ as $n \rightarrow \infty$.

Suppose that either one of the vectors v or w is zero, or that they are parallel. Since by continuity $|a|$ and $|b|$ are bounded by 3δ , necessarily we would have that the area of the parallelogram with side a_n and b_n tend to zero, as $n \rightarrow \infty$. In turn, this means that the area of a fundamental domain of \mathbb{R}^2 for the action of Γ_n tend to zero as $n \rightarrow \infty$. Applying Lemma 2.11 with $K = \overline{B_{3\delta}^{\mathbb{R}^2}(0)}$, by Proposition 3.1.4 in [AT04], the two dimensional Hausdorff measure of m_n tends to zero, thus contradicting Theorem 2.4. \square

3.3. Construction of the solution. By Corollary 2.10 and Lemma 3.1, up to extract a subsequence, the sequence (u_n) converges to a horoconvex function u , uniformly on any compact set.

Let $a, b \in \mathbb{R}^2$ given by Lemma 3.3, and define $\Gamma \subset \text{Iso}(\mathbb{R}^2)$ as the direct product of $\langle a \rangle$ and $\langle b \rangle$. Since a and b are linearly independent vectors, \mathbb{R}^2/Γ is a torus.

Lemma 3.4. *The function u is Γ -invariant.*

Proof. Let $y \in \mathbb{R}^2$ and $\gamma \in \Gamma$. Then, for every $\epsilon > 0$

$$\begin{aligned}
 |u(\gamma y) - u(y)| &= |u(y + ka + k'b) - u(y)| \\
 (7) \quad &\leq |u(y + ka + k'b) - u_n(y + ka + k'b)| \\
 (8) \quad &+ |u_n(y + ka + k'b) - u_n(y + ka_n + k'b_n)| \\
 (9) \quad &+ |u_n(y + ka_n + k'b_n) - u_n(y)| \\
 (10) \quad &+ |u_n(y) - u(y)| < \epsilon
 \end{aligned}$$

for n large enough. In fact $k(a - a_n) + k'(b - b_n) \rightarrow 0$ as $n \rightarrow \infty$, and as the u_n are equi-Lipschitz on a sufficiently large compact set the absolute value at line (8) is smaller than $\epsilon/4$ for n large enough. Moreover the absolute value at line (9) is zero for every n by the Γ_n -invariance of u_n , and the absolute value at lines (7) and (10) are smaller than $\epsilon/4$ for n large enough by the uniform convergence of the u_n . Since $\epsilon > 0$ is arbitrary, this concludes the proof. \square

3.4. Convergence of metrics. In the preceding section, we have constructed a pair (u, Γ) . It remains to check that the induced metric m_u on $(\mathbb{R}^2, d_u)/\Gamma$ is isometric to (T, m) . Basically, one has to check that if the sequence (u_n) converges, then the sequence of induced metric converges. In the remaining of this section we will prove the following result, that ends the proof of the theorem, because on compact metric spaces, uniform convergence imply Gromov–Hausdorff convergence, and the Gromov–Hausdorff limit is unique.

Proposition 3.5. *The sequence (m_n) uniformly converges to m_u .*

Lemma 3.6. *Let $K \subset \mathbb{R}^2$ be compact. Then*

$$E(K) = \text{closure}(\cup_{x \in K} \cup_n \{y | d_{u_n}(x, y) \leq \text{diam}\})$$

is compact.

Proof. We have

$$d_{\mathbb{H}^3} \left(\left(x, e^{-u_n(x)} \right), \left(y, e^{-u_n(y)} \right) \right) \leq d_{u_n}(x, y) \leq \text{diam} ,$$

but [Rat06, 4.6.1]

$$\cosh d_{\mathbb{H}^3} \left(\left(x, e^{-u_n(x)} \right), \left(y, e^{-u_n(y)} \right) \right) = 1 + \frac{\|x - y\|^2}{2e^{-u_n(x)}e^{-u_n(y)}}$$

so

$$\|x - y\| \leq \sqrt{2}e^{-c}\sqrt{\cosh(\text{diam}) - 1}$$

where c is the uniform lower bound of the u_n . The result follows because K is compact. \square

Corollary 3.7. *Let $K \subset \mathbb{R}^2$ be compact. Let $\mathcal{L}(K)$ be the set of shortest paths for any d_n between points of K (we don't ask the shortest path to be contained in K). Then there exists a constant α such that $\forall c \in \mathcal{L}(K)$, $L_{\mathbb{R}^2}(c) \leq \alpha$.*

Proof. By Lemma 2.11 applied to K , for any $x, y \in K$,

$$d_{u_n}(x, y) \leq \lambda_2(K) \text{diam}(K) .$$

Let c be a shortest path for u_n between x and y , so that $L_{u_n}(c) = d_{u_n}(x, y)$.

The shortest path is contained in the compact set $E(K)$ given by Lemma 3.6, so by Lemma 2.11 again, but applied to $E(K)$, $L_{\mathbb{R}^2}(c) \leq \frac{\lambda_2(K)}{\lambda_1(E(K))} \text{diam}(K)$. \square

Lemma 3.8. *Let $K \subset \mathbb{R}^2$ be compact. Then d_{u_n} uniformly converge to d_u on K .*

Proof. Let $\epsilon > 0$. As u_n uniformly converge to u , for n sufficiently large, $u \leq u_n + \epsilon$, so by Lemma 2.12

$$d_u \leq d_{u_n+\epsilon} + 4\epsilon .$$

Let $x, y \in K$ and let c be a shortest path for u_n between x and y . Then

$$d_{u_n+\epsilon}(x, y) \leq L_{u_n+\epsilon}(c)$$

and from (4), the fact that the u_n are uniformly bounded and Corollary 3.7, there exists a constant β depending only on K such that

$$L_{u_n+\epsilon}(c) \leq L_{u_n}(c) + |e^{2\epsilon} - 1|\beta(K)$$

and as $L_{u_n}(c) = d_{u_n}(x, y)$ we obtain

$$d_u(x, y) - d_{u_n}(x, y) \leq 4\epsilon + |e^{2\epsilon} - 1|\beta(K) .$$

Exchanging the role of u and u_n , we finally obtain

$$|d_u(x, y) - d_{u_n}(x, y)| \leq 4\epsilon + |e^{2\epsilon} - 1|\beta(K) .$$

\square

Remark 3.9. A result of A.D. Alexandrov gives a weaker convergence of the induced metrics for any convex surfaces converging in the Hausdorff sense to a convex surface in the hyperbolic space, see [Slu].

Let $\tilde{\varphi}_n$ be the linear isomorphism sending a to a_n and b to b_n . Hence clearly, for any $\gamma \in \Gamma$,

$$(11) \quad \tilde{\varphi}_n(\gamma \cdot x) = \tilde{\varphi}_n(\gamma) \cdot \tilde{\varphi}_n(x)$$

where γ is considered as a vector of \mathbb{R}^2 . The map $\tilde{\varphi}_n$ descends to a homeomorphism φ_n between \mathbb{R}^2/Γ and \mathbb{R}^2/Γ_n .

Lemma 3.10. *Let $K \subset \mathbb{R}^2$. Then on K ,*

$$x \mapsto d_{u_n}(\tilde{\varphi}_n(x), x)$$

uniformly converge to zero and $d_{u_n}(\tilde{\varphi}_n(x), \tilde{\varphi}_n(y))$ uniformly converge to $d_u(x, y)$.

Proof. $\tilde{\varphi}_n$ converge to the identity map, uniformly on any compact for the Euclidean metric. The first result follows from Lemma 2.11. The second result follows easily from the first one, the triangle inequality and Lemma 3.8. \square

Lemma 3.11. *There exists a compact set $K \subset \mathbb{R}^2$ such that for any n , for any p, q in the torus, a lift of a shortest path for m_n between p and q is contained in K .*

Proof. Let $K = E(D)$, where D is the compact set obtained in Lemma 3.1. By definition of D , there exists a lift x of p in D . By construction of K , the ball for d_n centred at x with radius $m_n(p, q)$ is contained in K . \square

Let C be the closure of $\cup_n \tilde{\varphi}_n(K)$, where K is given by Lemma 3.11. C is a compact set.

Lemma 3.12. *For any $\nu > 0$, if n is sufficiently large, for any $p, q \in T$, if x and y are respective lifts to the set C defined above, if $d_u(x, y) = m(p, q)$, then*

$$m_n(\varphi_n(p), \varphi_n(q)) \leq d_{u_n}(\tilde{\varphi}_n(x), \tilde{\varphi}_n(y)) \leq m_n(\varphi_n(p), \varphi_n(q)) + \nu.$$

Proof. As $d_u(x, y) = m(p, q)$, for any $\gamma \in \Gamma$,

$$d_u(x, \gamma \cdot y) \geq d_u(x, y).$$

By Lemma 3.10, uniformly on C , if n is sufficiently large,

$$d_{u_n}(\tilde{\varphi}_n(x), \tilde{\varphi}_n(\gamma) \cdot \tilde{\varphi}_n(y)) + \nu \geq d_{u_n}(\tilde{\varphi}_n(x), \tilde{\varphi}_n(y)).$$

The result follows because $m_n(\varphi_n(p), \varphi_n(q))$ is the minimum on Γ of all the $d_{u_n}(\tilde{\varphi}_n(x), \tilde{\varphi}_n(\gamma) \cdot \tilde{\varphi}_n(y))$. \square

Proof of Proposition 3.5. Let $\epsilon > 0$. For any $p, q \in T$, with the notations of Lemma 3.12, for n large enough,

$$\begin{aligned} & m_n(\varphi_n(p), \varphi_n(q)) - m(p, q) \\ & \leq d_{u_n}(\tilde{\varphi}_n(x), \tilde{\varphi}_n(y)) - d_u(x, y) \\ & \leq d_{u_n}(\tilde{\varphi}_n(x), x) + d_{u_n}(\tilde{\varphi}_n(x), y) + d_{u_n}(x, y) - d_u(x, y). \end{aligned}$$

Applying Lemma 3.8 and Lemma 3.10 on the compact set C , for n sufficiently large, the last quantity above is less than ϵ , independently of x and y . For the same reasons, the same conclusion holds for

$$m(p, q) - m_n(\varphi_n(p), \varphi_n(q))$$

$$\begin{aligned} &\leq d_u(x, y) - d_{u_n}(\tilde{\varphi}_n(x), \tilde{\varphi}_n(y)) + \nu \\ &\leq d_u(x, y) - d_{u_n}(x, y) + d_{u_n}(x, y) - d_{u_n}(\tilde{\varphi}_n(x), \tilde{\varphi}_n(y)) + \nu . \end{aligned}$$

□

REFERENCES

- [AKP] S. Alexander, V. Kapovitch, and A. Petrunin. *Alexandrov geometry*. In preparation. Draft version available.
- [AKP08] S. Alexander, V. Kapovitch, and A. Petrunin. An optimal lower curvature bound for convex hypersurfaces in Riemannian manifolds. *Illinois J. Math.*, 52(3):1031–1033, 2008.
- [Ale06] A. D. Alexandrov. *A. D. Alexandrov selected works. Part II*. Chapman & Hall/CRC, Boca Raton, FL, 2006. Intrinsic geometry of convex surfaces, Edited by S. S. Kutateladze, Translated from the Russian by S. Vakhrameyev.
- [AT04] L. Ambrosio and P. Tilli. *Topics on analysis in metric spaces*, volume 25 of *Oxford Lecture Series in Mathematics and its Applications*. Oxford University Press, Oxford, 2004.
- [BBI01] D. Burago, Y. Burago, and S. Ivanov. *A course in metric geometry*, volume 33 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2001.
- [BGP92] Yu. Burago, M. Gromov, and G. Perelman. A. D. Aleksandrov spaces with curvatures bounded below. *Uspekhi Mat. Nauk*, 47(2(284)):3–51, 222, 1992.
- [BH37] W. Blaschke and G. Herglotz. Über die Verwirklichung einer geschlossenen Fläche mit vorgeschriebenem Bogenelement im Euklidischen Raum. *Sitzungsber. Bayer. Akad. Wiss., Math.-Naturwiss. Abt.*, (2):229–230, 1937.
- [BH99] M. Bridson and A. Haefliger. *Metric spaces of non-positive curvature*, volume 319 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1999.
- [BI08] A. Bobenko and I. Izvestiev. Alexandrov’s theorem, weighted Delaunay triangulations, and mixed volumes. *Ann. Inst. Fourier (Grenoble)*, 58(2):447–505, 2008.
- [CEG06] R. D. Canary, D. B. A. Epstein, and P. L. Green. Notes on notes of Thurston [mr0903850]. In *Fundamentals of hyperbolic geometry: selected expositions*, volume 328 of *London Math. Soc. Lecture Note Ser.*, pages 1–115. Cambridge Univ. Press, Cambridge, 2006. With a new foreword by Canary.
- [CX15] J.-E. Chang and L. Xiao. The Weyl problem with nonnegative Gauss curvature in hyperbolic space. *Canad. J. Math.*, 67(1):107–131, 2015.
- [FI09] F. Fillastre and I. Izvestiev. Hyperbolic cusps with convex polyhedral boundary. *Geom. Topol.*, 13(1):457–492, 2009.
- [Fil07] F. Fillastre. Polyhedral realisation of hyperbolic metrics with conical singularities on compact surfaces. *Ann. Inst. Fourier (Grenoble)*, 57(1):163–195, 2007.
- [Gro86] M. Gromov. *Partial differential relations*, volume 9 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1986.
- [GSS09] B. Guan, J. Spruck, and M. Szapiel. Hypersurfaces of constant curvature in hyperbolic space. I. *J. Geom. Anal.*, 19(4):772–795, 2009.
- [IRV15] J. Itoh, J. Rouyer, and C. Vilcu. Moderate smoothness of most Alexandrov surfaces. *Internat. J. Math.*, 26(4):1540004 (14 pages), 2015.
- [Izm08] I. Izvestiev. A variational proof of Alexandrov’s convex cap theorem. *Discrete Comput. Geom.*, 40(4):561–585, 2008.
- [Izm13] I. Izvestiev. Infinitesimal rigidity of smooth convex surfaces through the second derivative of the Hilbert-Einstein functional. *Dissertationes Math. (Rozprawy Mat.)*, 492:58, 2013.
- [Lab92] F. Labourie. Métriques prescrites sur le bord des variétés hyperboliques de dimension 3. *J. Differential Geom.*, 35(3):609–626, 1992.

- [Pog73] A. V. Pogorelov. *Extrinsic geometry of convex surfaces*. American Mathematical Society, Providence, R.I., 1973. Translated from the Russian by Israel Program for Scientific Translations, Translations of Mathematical Monographs, Vol. 35.
- [Rat06] J. Ratcliffe. *Foundations of hyperbolic manifolds*, volume 149 of *Graduate Texts in Mathematics*. Springer, New York, second edition, 2006.
- [Ric12] T. Richard. *Ricci flow without upper bounds on the curvature and the geometry of some metric spaces*. Theses, Université de Grenoble, September 2012.
- [Roc97] R. Rockafellar. *Convex analysis*. Princeton Landmarks in Mathematics. Princeton University Press, Princeton, NJ, 1997. Reprint of the 1970 original, Princeton Paperbacks.
- [RS94] H. Rosenberg and J. Spruck. On the existence of convex hypersurfaces of constant Gauss curvature in hyperbolic space. *J. Differential Geom.*, 40(2):379–409, 1994.
- [RW98] T. Rockafellar and R. Wets. *Variational analysis*, volume 317 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1998.
- [Sch06] J.-M. Schlenker. Hyperbolic manifolds with convex boundary. *Invent. Math.*, 163(1):109–169, 2006.
- [Slu] D. Slutskiy. Compact domains with prescribed convex boundary metrics in quasi-Fuchsian manifolds. arXiv:1405.1650.
- [Slu14] D. Slutskiy. Polyhedral metrics on the boundaries of convex compact quasi-Fuchsian manifolds. *C. R. Math. Acad. Sci. Paris*, 352(10):831–834, 2014.

UNIVERSITÉ DE CERGY-PONTOISE, UMR CNRS 8088, F-95000 CERGY-PONTOISE, FRANCE

E-mail address: `francois.fillastre@u-cergy.fr`

DEPARTMENT OF MATHEMATICS, FREE UNIVERSITY OF BERLIN, ARNIMALLEE 2, D-14195 BERLIN, GERMANY

E-mail address: `izmestiev@math.fu-berlin.de`

UNIVERSITÉ PARIS 13, SORBONNE PARIS CITÉ, LAGA, CNRS (UMR 7539), 99 AV. JEAN-BAPTISTE CLÉMENT, 93430 VILLETANEUSE, FRANCE

E-mail address: `veronelli@math.univ-paris13.fr`